Your Guide to Understanding Genetic Conditions

PSAP gene

prosaposin

Normal Function

The *PSAP* gene provides instructions for making a protein called prosaposin. This protein is involved in a number of biological functions, including the development of the nervous system and the reproductive system. Prosaposin is the precursor of four smaller proteins called saposin A, B, C, and D, which are produced when prosaposin is broken up (cleaved).

The individual saposins are found in cellular structures called lysosomes, which are the cell's recycling centers. The saposins help lysosomal enzymes break down fatty substances called sphingolipids.

The saposin B protein works with several enzymes to break down sphingolipids. Its most critical biological role seems to be associated with the enzyme arylsulfatase A. This enzyme is involved in breaking down a subgroup of sphingolipids called sulfatides, especially in the nervous system's white matter, which consists of nerve fibers covered by myelin. Myelin is a substance that insulates and protects nerves. Saposin B may also play a role in transporting lipids to the outer surface of the cell so they can be recognized by the immune system.

The saposin C protein works with the enzyme beta-glucocerebrosidase to break down another sphingolipid called glucocerebroside. Saposins A and D are also involved in processing sphingolipids.

Health Conditions Related to Genetic Changes

metachromatic leukodystrophy

In a small number of individuals with metachromatic leukodystrophy, a disorder that causes deterioration of nervous system functions, researchers have identified *PSAP* gene mutations that result in a shortage (deficiency) of the saposin B protein. This deficiency interferes with the breakdown of sulfatides. As a result, these substances can accumulate to toxic levels in the nervous system.

The buildup of sulfatides gradually destroys myelin, the covering that protects nerves and promotes the efficient transmission of nerve impulses. Destruction of myelin leads to a loss of white matter (leukodystrophy) and impairment of nervous system function, resulting in the signs and symptoms of metachromatic leukodystrophy.

other disorders


In a few individuals, mutations in the *PSAP* gene interfere with the function of the saposin C protein, resulting in a disorder that resembles a severe form of Gaucher disease. Signs and symptoms of this condition include neurological problems and abnormal enlargement of the liver and spleen (hepatosplenomegaly). Without adequate saposin C activator protein, the glucocerebrosidase enzyme cannot break down glucocerebroside effectively. As a result, glucocerebroside accumulates in the body's tissues as it does in the classic form of Gaucher disease. A few *PSAP* gene mutations have also been identified in individuals with signs and symptoms resembling another leukodystrophy called Krabbe disease.

In addition, a few mutations in the *PSAP* gene have been identified that prevent the production of more than one of the saposin proteins. Individuals with these mutations have massive accumulation of sphingolipids in their nervous system and other organs. This accumulation results in very severe neurological disease, respiratory problems, and hepatosplenomegaly.

Chromosomal Location

Cytogenetic Location: 10q22.1, which is the long (q) arm of chromosome 10 at position 22.1

Molecular Location: base pairs 71,816,283 to 71,851,375 on chromosome 10 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- Prosaposin (sphingolipid activator protein-1)
- prosaposin (variant Gaucher disease and variant metachromatic leukodystrophy)
- prosaptides
- SAP1
- SAP2 (sphingolipid activator protein-2)

- SAP HUMAN
- SGP-1 (sulfoglycoprotein-1)

Additional Information & Resources

Educational Resources

- Basic Neurochemistry (sixth edition, 1999): Lysosomal Disease https://www.ncbi.nlm.nih.gov/books/NBK28215/
- Essentials of Glycobiology (first edition, 1999): Glycosphingolipid Degradation https://www.ncbi.nlm.nih.gov/books/NBK20729/#A1383

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28PSAP%5BTIAB%5D%29+OR+%28prosaposin%5BTIAB%5D%29+OR+%28%28SAP1%5BTIAB%5D%29+OR+%28prosaposin%5BTIAB%5D%29+OR+%28GLBA%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D

OMIM

- COMBINED SAPOSIN DEFICIENCY http://omim.org/entry/611721
- GAUCHER DISEASE, ATYPICAL, DUE TO SAPOSIN C DEFICIENCY http://omim.org/entry/610539
- KRABBE DISEASE, ATYPICAL, DUE TO SAPOSIN A DEFICIENCY http://omim.org/entry/611722
- PROSAPOSIN http://omim.org/entry/176801

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology http://atlasgeneticsoncology.org/Genes/PSAPID42980ch10q22.html
- ClinVar https://www.ncbi.nlm.nih.gov/clinvar?term=PSAP%5Bgene%5D
- HGNC Gene Family: Endogenous ligands http://www.genenames.org/cgi-bin/genefamilies/set/542

- HGNC Gene Symbol Report http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/ hgnc_data.php&hgnc_id=9498
- NCBI Gene https://www.ncbi.nlm.nih.gov/gene/5660
- UniProt http://www.uniprot.org/uniprot/P07602

Sources for This Summary

- Al-Hassnan ZN, Al Dhalaan H, Patay Z, Faqeih E, Al-Owain M, Al-Duraihem A, Faiyaz-Ul-Haque M. Sphingolipid activator protein B deficiency: report of 9 Saudi patients and review of the literature. J Child Neurol. 2009 Dec;24(12):1513-9. doi: 10.1177/0883073809341269. Review. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/19955343
- Basic Neurochemistry (sixth edition, 1999): Lysosomal Disease https://www.ncbi.nlm.nih.gov/books/NBK28215/
- Deconinck N, Messaaoui A, Ziereisen F, Kadhim H, Sznajer Y, Pelc K, Nassogne MC, Vanier MT, Dan B. Metachromatic leukodystrophy without arylsulfatase A deficiency: a new case of saposin-B deficiency. Eur J Paediatr Neurol. 2008 Jan;12(1):46-50. Epub 2007 Jul 5.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/17616409
- Diaz-Font A, Cormand B, Santamaria R, Vilageliu L, Grinberg D, Chabás A. A mutation within the saposin D domain in a Gaucher disease patient with normal glucocerebrosidase activity. Hum Genet. 2005 Jul;117(2-3):275-7. Epub 2005 Apr 23.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/15856305
- Essentials of Glycobiology (first edition, 1999): Glycosphingolipid Degradation https://www.ncbi.nlm.nih.gov/books/NBK20729/#A1383
- OMIM: GAUCHER DISEASE, ATYPICAL, DUE TO SAPOSIN C DEFICIENCY http://omim.org/entry/610539
- Grossi S, Regis S, Rosano C, Corsolini F, Uziel G, Sessa M, Di Rocco M, Parenti G, Deodato F, Leuzzi V, Biancheri R, Filocamo M. Molecular analysis of ARSA and PSAP genes in twenty-one Italian patients with metachromatic leukodystrophy: identification and functional characterization of 11 novel ARSA alleles. Hum Mutat. 2008 Nov;29(11):E220-30. doi: 10.1002/humu.20851. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/18693274
- OMIM: PROSAPOSIN http://omim.org/entry/176801
- Szymanska K, Lugowska A, Laure-Kamionowska M, Bekiesinska-Figatowska M, Gieruszczak-Bialek D, Musielak M, Eichler S, Giese AK, Rolfs A. Diagnostic difficulties in Krabbe disease: a report of two cases and review of literature. Folia Neuropathol. 2012;50(4):346-56. Review. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/23319190
- Tamargo RJ, Velayati A, Goldin E, Sidransky E. The role of saposin C in Gaucher disease. Mol Genet Metab. 2012 Jul;106(3):257-63. doi: 10.1016/j.ymgme.2012.04.024. Epub 2012 May 5. Review.
 - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/22652185
 Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3534739/

Reprinted from Genetics Home Reference: https://ghr.nlm.nih.gov/gene/PSAP

Reviewed: February 2013 Published: March 21, 2017

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services